

KELLER'S CONJECTURE ON THE EXISTENCE OF COLUMNS IN CUBE TILINGS OF $\mathbb{R}^n$

Magdalena Łysakowska and Krzysztof Przesański

Wydział Matematyki, Informatyki i Ekonometrii, Uniwersytet Zielonogórski

ul. Z. Szafrana 4a, 65-516 Zielona Góra, Poland

M.Lysakowska@wmie.uz.zgora.pl

K.Przeslawski@wmie.uz.zgora.pl

Abstract

It is shown that if $n \leq 6$, then each tiling of $\mathbb{R}^n$ by translates of the unit cube $[0, 1)^n$ contains a column; that is, a family of the form $\{[0, 1)^n + (s + ke_i) : k \in \mathbb{Z}\}$, where $s \in \mathbb{R}^n$ and e_i is an element of the standard basis of $\mathbb{R}^n$.

Key words: cube tiling, column.

1 Introduction

In his book [6], which appeared in 1907, Hermann Minkowski proved that every lattice tiling of $\mathbb{R}^n$ by unit cubes contains two cubes that have a common $(n - 1)$ -dimensional face, whenever $n \leq 3$. This readily implies that there is a column of unit cubes contained in the tiling. On the other hand, he conjectured that the same phenomenon holds in all dimensions. In 1930, Otto Heinrich Keller [2] extended Minkowski's conjecture to arbitrary cube tilings of $\mathbb{R}^n$. In fact, we have now two conjectures: the stronger, stated by Keller, which reads that each cube tiling of $\mathbb{R}^n$ contains a column and the weaker which reads that each cube tiling of $\mathbb{R}^n$ contains two cubes that share an $(n - 1)$ -dimensional face. In 1937, Keller published a short paper [3] where he claimed that he proved his conjecture for $n \leq 6$. He also expressed a supposition that the conjecture is not valid in dimensions greater than 6. There are no rigorous proofs in the paper however. In 1940, Oskar Perron [7] published a complete proof of the weaker conjecture for dimensions not exceeding 6. He left aside the original conjecture of Keller. Apparently, he

was focused on Minkowski's conjecture, which was verified by Hajós [1] only one year later. It was Perron who popularized the weaker conjecture under the name of Keller. In 1992, Jeff Lagarias and Peter Shor [4] discovered a counterexample to the weaker conjecture in dimension 10. Ten years later John Mackey [5] found a counterexample in dimension 8. This implies that both conjectures, the weaker and the stronger, are not valid in any dimension greater than 7. For dimension 7, both problems are completely open. In the present note we show that the original Keller's conjecture is valid in all dimensions up to 6. Our proof is based on Perron's approach. However, certain substantial modifications of his method were necessary: He concentrated his attention on local configurations of cubes, whereas we have to play with all cubes of tiling. This forces us to work with an appropriate enumeration of cubes.

2 The existence of columns

We define a *cube* in the n -dimensional Euclidean space $\mathbb{R}^n$ to be any translate of the unit cube $[0, 1]^n$. Let T be a subset of $\mathbb{R}^n$. The family $[0, 1]^n + T := \{[0, 1]^n + t : t \in T\}$ is said to be a *cube tiling* of $\mathbb{R}^n$ if for each pair of distinct vectors $s, t \in T$ the cubes $[0, 1]^n + s$ and $[0, 1]^n + t$ are disjoint and $\bigcup [0, 1]^n + T = \mathbb{R}^n$. We refer to T as a set that determines a cube tiling.

As usual, we denote by $\mathbb{Z}$ the set of all integers while the set of positive integers is denoted by $\mathbb{N}$. Let $n \in \mathbb{N}$. The set $\{1, 2, \dots, n\}$ is denoted by $[n]$.

Let us recall the following fundamental result from [2].

Theorem 1 (O. H. Keller, 1930) *If $[0, 1]^n + T$ is a cube tiling of $\mathbb{R}^n$, then for each pair of distinct elements $s, t \in T$ there is $j \in [n]$ such that $|s_j - t_j| \in \mathbb{N}$.*

Proof. Suppose the theorem is not valid. Then there is a set T that determines a cube tiling of $\mathbb{R}^n$ which contains a pair of distinct elements $s, t \in T$ such that $s_j - t_j \notin \mathbb{Z} \setminus \{0\}$ for each $j \in [n]$. Clearly, the set $S = T - t$ determines a cube tiling as well. Let $u = s - t$. The elements u and 0 belong to S and $u_j \notin \mathbb{Z} \setminus \{0\}$ for each $j \in [n]$. For $x \in \mathbb{R}^n$, let $i(x) := \max\{i : |x_j| < 1 \text{ for } j \leq i\}$, where we have assumed that $\max \emptyset = 0$. We have $i(u) < n$, otherwise cubes $[0, 1]^n$ and $[0, 1]^n + u$ would intersect contradicting the assumption that S determines a cube tiling. Let $k := i(u) + 1$ and let $V := \{v \in S : v_k - u_k \in \mathbb{Z}\}$. If ℓ is a straight line intersecting one of the cubes $[0, 1]^n + v$, $v \in V$, which in addition is parallel to the k -th coordinate axis, then $\ell \subset \bigcup_{v \in V} ([0, 1]^n + v)$. This observation leads to the conclusion that the set $U := (V - \lfloor u_k \rfloor e_k) \cup (S \setminus V)$ determines a cube tiling.

Obviously, $r := u - \lfloor u_k \rfloor e_k$ and 0 are elements of U . Moreover, $i(r) = i(u) + 1$ and $r_j \notin \mathbb{Z} \setminus \{0\}$ for every $j \in [n]$. Now, we can replace S by U , and continue in this manner eventually arriving to a set that determines a cube tiling and contains 0 and an element w such that $i(w) = n$ and $w_j \notin \mathbb{Z} \setminus \{0\}$ for every $j \in [n]$ which, as we know, is impossible. $\square$

Let us suppose that for each $j \in [n]$ a mapping $\varepsilon_j: \mathbb{R} \rightarrow \mathbb{N}$ is given such that for each element $x \in \mathbb{R}$ the restriction $\varepsilon_j|x + \mathbb{Z}$ is a bijection between the sets $x + \mathbb{Z} := \{x + k: k \in \mathbb{Z}\}$ and $\mathbb{N}$. The mapping $\varepsilon: \mathbb{R}^n \rightarrow \mathbb{N}^n$ defined by the formula

$$\varepsilon(x) = \varepsilon(x_1, \dots, x_n) = (\varepsilon_1(x_1), \dots, \varepsilon_n(x_n))$$

is said to be a *natural code* (of $\mathbb{R}^n$). The vector $\varepsilon(x)$ is referred to as the code of x .

Theorem 2 *Fix a natural code $\varepsilon: \mathbb{R}^n \rightarrow \mathbb{N}^n$. Then a set $T \subseteq \mathbb{R}^n$ determines a cube tiling of $\mathbb{R}^n$ if and only if $\varepsilon(T) = \mathbb{N}^n$ and for every pair of distinct elements $s, t \in T$ there is $j \in [n]$ such that $|s_j - t_j| \in \mathbb{N}$.*

Proof. ($\Rightarrow$) By Keller's theorem, it suffices to show that $\varepsilon(T) = \mathbb{N}^n$. We proceed by induction with respect n . For $n = 1$ the assertion is a consequence of the definition of a natural code and the fact that each set determining a cube tiling of $\mathbb{R}^1$ coincides with one of the cosets $\mathbb{Z} + x$, $x \in \mathbb{R}$. Fix $k \in \mathbb{N}$, and define the set

$$T^k := \{t = (t_1, \dots, t_n) \in T: \varepsilon_n(t_n) = k\}.$$

If $n > 1$, then $\mathbb{R}^{n-1}$ is a non-trivial Euclidean space. Let $T_{n'}^k \subset \mathbb{R}^{n-1}$ be the image of T^k under the projection $x = (x_1, \dots, x_n) \mapsto x_{n'} := (x_1, \dots, x_{n-1})$ of $\mathbb{R}^n$ onto $\mathbb{R}^{n-1}$. Let us show now that $T_{n'}^k$ determines a cube tiling of $\mathbb{R}^{n-1}$. Let $x \in \mathbb{R}^{n-1}$ and let $T(x) = \{t \in T: ([0, 1]^n + t) \cap (\{x\} \times \mathbb{R}) \neq \emptyset\}$. Since $[0, 1]^n + T$ is a cube tiling of $\mathbb{R}^n$, the set $T(x) + [0, 1]^n$ is a covering of the set $\{x\} \times \mathbb{R}$ by disjoint sets. Therefore, $T(x)_n = \{t_n: t \in T(x)\}$ determines a cube tiling of $\mathbb{R}$. The latter set is transformed by ε_n bijectively onto $\mathbb{N}$. In particular, there is an element $t \in T(x)$ such that $\varepsilon_n(t_n) = k$; equivalently, $t \in T^k$ and $x \in [0, 1]^{n-1} + t_{n'}$. Since x is arbitrary, we deduce that $T_{n'}^k$ determines a cube tiling of $\mathbb{R}^{n-1}$, as announced. Let $\varepsilon_{n'} := (\varepsilon_1, \dots, \varepsilon_{n-1})$. By the induction hypothesis, $\varepsilon_{n'}(T_{n'}^k) = \mathbb{N}^{n-1}$. Therefore, $\varepsilon(T^k) = \mathbb{N}^{n-1} \times \{k\}$. Finally,

$$\varepsilon(T) = \bigcup_{k \in \mathbb{N}} \varepsilon(T^k) = \bigcup_{k \in \mathbb{N}} \mathbb{N}^{n-1} \times \{k\} = \mathbb{N}^n.$$

($\Leftarrow$) As ε maps T ‘onto’ $\mathbb{N}^n$, it maps T^k ‘onto’ $\mathbb{N}^{n-1} \times \{k\}$. Hence $\varepsilon_{n'}$ maps $T_{n'}^k$ ‘onto’ $\mathbb{N}^{n-1}$. Moreover, by our assumptions for every two elements $s, t \in T_{n'}^k$ there is $j \in [n-1]$ such that $|s_j - t_j| \in \mathbb{N}$. Therefore, $T_{n'}^k$ determines a cube tiling of $\mathbb{R}^{n-1}$ by the induction hypothesis. Let $x \in \mathbb{R}^{n-1}$. Then for each number $k \in \mathbb{N}$ there is $t^k \in T^k$ such that $x \in [0, 1)^{n-1} + t_{n'}^k$. Since for every pair of distinct elements $r, s \in \{t^k : k \in \mathbb{N}\}$ there is $j \in [n]$ such that $|r_j - s_j| \in \mathbb{N}$ and the cubes $[0, 1)^{n-1} + t_{n'}^k$, $k \in \mathbb{N}$, intersect, the set $\{t_{n'}^k : k \in \mathbb{N}\}$ determines a cube tiling of $\mathbb{R}$. Thus, $\{x\} \times \mathbb{R} \subseteq \bigcup_{k \in \mathbb{N}} ([0, 1)^n + t^k) \subseteq \bigcup ([0, 1)^n + T)$. Consequently, $\bigcup ([0, 1)^n + T) = \mathbb{R}^n$. $\square$

Two vectors x and $y \in \mathbb{R}^n$ are $\mathbb{Z}$ -*distinguishable* if there is $j \in [n]$ such that $x_j \neq y_j$ and $y_j \in x_j + \mathbb{Z}$. A system of vectors is called $\mathbb{Z}$ -distinguishable, or shortly distinguishable, if any two vectors of this system are $\mathbb{Z}$ -distinguishable.

Let $l < n$. A family of boxes F is said to be an l -*column* if there is a set of vectors $S \subset \mathbb{R}^n$ such that the following conditions are satisfied:

- (1) $F = S + [0, 1)^n$;
- (2) there is $i \in [n]$ such that the mapping $x \mapsto x_i$ transforms S bijectively into a set that determines a cube partition of $\mathbb{R}^1$;
- (3) there are l indices $j \in [n]$ such that the sets $S_j := \{x_j : x \in S\}$ are singletons.

We refer to S as a set that determines an l -column. Every $(n-1)$ -column contained in $\mathbb{R}^n$ is called in short a *column*.

Two sets of vectors $F, G \subseteq \mathbb{R}^n$ are *isomorphic*, if there are a bijection $f: F \rightarrow G$ and a permutation $\sigma: [n] \rightarrow [n]$ such that for every pair of vectors $x, y \in F$ and for each $j \in [n]$ the following conditions are satisfied:

- (1) $x_j = y_j$ if and only if $f(x)_{\sigma(j)} = f(y)_{\sigma(j)}$,
- (2) $|x_j - y_j| \in \mathbb{N}$ if and only if $|f(x)_{\sigma(j)} - f(y)_{\sigma(j)}| \in \mathbb{N}$.

Let x be an element of $\mathbb{R}^n$. In what follows, we often write $x : x_1 \dots x_n$ instead of $x = (x_1, \dots, x_n)$, according to the convention adopted by Perron [7]. The coordinates of the vectors belonging to any set that determines a cube tiling of $\mathbb{R}^n$ are denoted by Roman or Greek lower case letters. We apply somewhat untypical convention that if various coordinates of a vector are represented by the same letter (possibly with the same upper and lower indices), then it does not imply that they have the same value. Starting from the proof of Theorem 6, it is tacitly assumed that when we talk about

the set of vectors that determines a cube tiling, a natural code of $\mathbb{R}^n$ is already defined. Such a code serves as a system of coordinates of the cube tiling. Lower indices of coordinates of a vector correspond to the code of this vector; e.g. the string of symbols $w : a_1\alpha_3^5a_2$ means that the vector $w = (a_1, \alpha_3^5, a_2)$ has the code $\varepsilon(w) = (1, 3, 2)$. If we have two vectors whose i -th coordinates are denoted by the same lower case letter with the same upper index, if there is any, then they differ by an integer; e.g. if $w_1 : a_1a_1\alpha_1^2$ and $w_2 : a_2b_1\alpha_3^2$, then $(w_1)_1 - (w_2)_1 = a_1 - a_2$ and $(w_1)_3 - (w_2)_3 = \alpha_1^2 - \alpha_3^2$ are non-zero integers. If i -th coordinates of two vectors are denoted by different Roman letters, then their difference is not an integer; e.g. if w_1 and w_2 are as above, then $(w_1)_2 = a_1$ while $(w_2)_2 = b_1$, therefore, $(w_1)_2 - (w_2)_2 = a_1 - b_1$ is not an integer. As is seen from the examples, Roman letters occur only with lower indices while Greek letters are equipped with lower and upper indices. The value of a coordinate of a vector is denoted by a Greek letter when it is not explicit; e.g. the third coordinates of the vectors $w_1 : a_1a_2\alpha_2^3$ and $w_2 : a_1a_1\alpha_2^4$ have the same code but it is not decided whether they are equal or different. There is only one place, the proof of Lemma 1, where coordinates of a vector are denoted by Roman capital letters. Their use will be explained therein.

Theorem 3 *Let n be a positive integer. If every cube tiling of $\mathbb{R}^n$ contains a column, then every cube tiling of $\mathbb{R}^m$ contains an $(n - 1)$ -column, for each $m > n$.*

Theorem 4 *Every cube tiling of $\mathbb{R}^2$ contains a column.*

Proof. Let T be an arbitrary set which determines a cube tiling of $\mathbb{R}^2$ and let $\varepsilon : \mathbb{R}^2 \rightarrow \mathbb{N}^2$ be a natural code. Let us consider these elements of T which have the codes $(k, 1)$, $k \in \mathbb{N}$. By Theorem 2 and the notation introduced above, they can be written as follows

$$w_{1,1} : a_1 \quad a_1,$$

$$w_{1,l} : a_l \quad \alpha_1^{l-1}, \quad l \geq 2.$$

If $\alpha_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,k}$, $k \geq 1$, determine a column. Let us suppose now that at least one of the numbers α_1^i , $i \geq 1$, is distinct from a_1 . We can assume that it is α_1^1 , as if $\alpha_1^1 = a_1$ and for example $\alpha_1^5 \neq a_1$, then we can change our natural code replacing ε_1 by the composite $\tau \circ \varepsilon_1$, where τ is the transposition (26), arriving in this way to a system of vectors where $w_{1,2}$ has its second coordinate different from a_1 . Now, let us take into account all vectors belonging to T with codes $(1, l)$, $l \geq 2$. As these

vectors together with $w_{1,1}$ and $w_{1,2}$ are $\mathbb{Z}$ -distinguishable, they must have the following form

$$w_{2,l-1} : \begin{matrix} a_1 & a_l \end{matrix}, \quad l \geq 2.$$

Thus, they together with $w_{1,1}$ determine a column. $\square$

Theorem 5 *Every cube tiling of $\mathbb{R}^3$ contains a column.*

Proof. Let T be an arbitrary set which determines a cube tiling of $\mathbb{R}^3$ and let $\varepsilon: \mathbb{R}^3 \rightarrow \mathbb{N}^3$ be a natural code. By Theorems 3 and 4, the set T contains vectors which determine a 1-column. Passing to an isomorphic system if necessary, we can assume that the vectors determining our 1-column are as follows

$$w_{1,1} : \begin{matrix} a_1 & a_1 & a_1 \end{matrix},$$

$$w_{1,l} : \begin{matrix} a_l & a_1 & \alpha_1^{l-1} \end{matrix}, \quad l \geq 2.$$

If $\alpha_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,k}$, $k \geq 1$, determine a column. Suppose that $\alpha_1^i \neq a_1$ for some $i \geq 1$. As previously, we can assume that $\alpha_1^1 \neq a_1$. Consider all vectors belonging to T with codes $(1,1,l)$, $l \geq 2$. As these vectors together with $w_{1,1}$ and $w_{1,2}$ are $\mathbb{Z}$ -distinguishable, they can be written in the following form

$$w_{2,l-1} : \begin{matrix} a_1 & \beta_1^{l-1} & a_l \end{matrix}, \quad l \geq 2.$$

If $\beta_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,1}$, $w_{2,k}$, $k \geq 1$, determine a column. Suppose that at least one of the coordinates β_1^i , $i \geq 1$, is different from a_1 . By the same reason as in the preceding proof, we can assume that $\beta_1^1 \neq a_1$. Now, let us consider all vectors from T with codes $(1,l,2)$, $l \geq 2$. As these vectors together with $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$ are $\mathbb{Z}$ -distinguishable, they must have the following form

$$w_{3,l-1} : \begin{matrix} a_1 & \beta_l^1 & a_2 \end{matrix}, \quad l \geq 2.$$

Thus, they and $w_{2,1}$ determine a column. $\square$

Theorem 6 *Every cube tiling of $\mathbb{R}^4$ contains a column.*

Proof. Let T be an arbitrary set which determines a cube tiling of $\mathbb{R}^4$. By Theorems 3 and 5, the set T contains vectors which determine a 2-column. Passing to an isomorphic system if necessary, we can assume that the vectors determining our 2-column are as follows

$$w_{1,1} : \begin{matrix} a_1 & a_1 & a_1 & a_1 \end{matrix},$$

$$w_{1,l} : \begin{matrix} a_l & a_1 & a_1 & \alpha_1^{l-1} \end{matrix}, \quad l \geq 2.$$

If $\alpha_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,k}$, $k \geq 1$, determine a column. Suppose that $\alpha_1^i \neq a_1$ for some $i \geq 1$. We can assume that $\alpha_1^1 \neq a_1$. Now, let us consider all vectors from T with codes $(1, *, *, l)$, $l \geq 2$, where $*$ can take any value from $\mathbb{N}$, such that their second and third coordinates are different from a_l , $l \geq 2$. (Such vectors exist, e.g. the vector whose code is $(1, 1, 1, 5)$ has the required property.) Let us pick a vector from among them whose *middle* coordinates (second and third) differ from a_1 at as many places as possible. Let us change the natural code so that the vectors $w_{1,l}$, $l \geq 1$ remain unaffected while the picked vector has its code equal to $(1, 1, 1, 2)$. By $\mathbb{Z}$ -distinguishability of this vector from $w_{1,1}$ and $w_{1,2}$, it can be written as follows

$$w_{2,1} : \quad a_1 \quad \beta_1^1 \quad \beta_1^2 \quad a_2.$$

Three cases have to be considered:

Case 1. $\beta_1^1 = \beta_1^2 = a_1$.

Then take all the vectors with codes $(1, 1, 1, l)$, $l \geq 3$. Since each of them must be distinguishable from the vectors $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, we deduce that they can be written in the form

$$w_{2,l-1}'' : \quad a_1 \quad \beta_1^{2l-3} \quad \beta_1^{2l-2} \quad a_l, \quad l \geq 3.$$

As $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\beta_1^i = a_1$ for all $i \geq 3$. Thus, the vectors $w_{1,1}$, $w_{2,1}$, $w_{2,l}''$, $l \geq 2$, determine a column.

Case 2. Exactly one of the coordinates β_1^1 and β_1^2 is equal to a_1 .

Then we can assume that $\beta_1^2 = a_1$, as in the other case we would change the order of the second and the third coordinate. Consider all vectors with codes $(1, l, 1, 2)$, $l \geq 2$. By the distinguishability, they can be written in the form

$$w_{2,l}' : \quad a_1 \quad \beta_l^1 \quad \gamma_1^{l-1} \quad a_2, \quad l \geq 2.$$

As $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\gamma_1^i = a_1$ for all $i \geq 1$. Thus, the vectors $w_{2,1}$ and $w_{2,l}'$, $l \geq 2$, determine a column.

Case 3. Both coordinates β_1^1 and β_1^2 are different from a_1 .

Then consider all vectors with codes $(1, l, 1, 2)$ and $(1, 1, l, 2)$, $l \geq 2$. By their distinguishability from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they have the form

$$w_{2,l} : \quad a_1 \quad \beta_l^1 \quad \gamma_1^{l-1} \quad a_2,$$

$$w_{3,l-1} : \quad a_1 \quad \delta_1^{l-1} \quad \beta_l^2 \quad a_2, \quad l \geq 2.$$

Since the vectors $w_{2,l}$, $l \geq 2$, and $w_{3,k}$, $k \geq 1$, are $\mathbb{Z}$ -distinguishable, we have $\delta_1^i = \beta_1^1$ for all $i \geq 1$ or $\gamma_1^i = \beta_1^2$ for all $i \geq 1$. If the first case takes place the vectors $w_{2,1}$, $w_{3,k}$, $k \geq 1$, determine a column; otherwise the vectors $w_{2,l}$, $l \geq 1$, determine a column. $\square$

Theorem 7 *Every cube tiling of $\mathbb{R}^5$ contains a column.*

Proof. Let T be an arbitrary set which determines a cube tiling of $\mathbb{R}^5$. By Theorems 3 and 6, the set T contains vectors which determine a 3-column. Passing to an isomorphic system if necessary, we can suppose that the vectors determining our 3-column are as follows

$$w_{1,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{1,l} : a_l \ a_1 \ a_1 \ a_1 \ \alpha_1^{l-1}, \quad l \geq 2.$$

If $\alpha_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,k}$, $k \geq 1$, determine a column. Suppose that $\alpha_1^i \neq a_1$ for some $i \geq 1$. As before, we can assume that $\alpha_1^1 \neq a_1$. Now, let us consider all vectors from T with codes $(1, *, *, *, l)$, $l \geq 2$, where $*$ can take any value from $\mathbb{N}$, such that their middle coordinates (second, third and fourth) are different from a_l , $l \geq 2$. Let us pick a vector from among them whose middle coordinates differ from a_1 at as many places as possible. Similarly as in the proof of Theorem 6, we can assume that this vector has its code equal to $(1, 1, 1, 1, 2)$. By its distinguishability from $w_{1,1}$ and $w_{1,2}$, it can be written in the form

$$w_{2,1} : a_1 \ \beta_1^1 \ \beta_1^2 \ \beta_1^3 \ a_2.$$

Four cases have to be considered:

Case 1. $\beta_1^1 = \beta_1^2 = \beta_1^3 = a_1$.

Then take all vectors with codes $(1, 1, 1, 1, l)$, $l \geq 3$. By the distinguishability, these vectors have the form

$$w_{2,l-1}'' : a_1 \ \beta_1^{3l-5} \ \beta_1^{3l-4} \ \beta_1^{3l-3} \ a_l, \quad l \geq 3.$$

As $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\beta_1^i = a_1$ for all $i \geq 4$. Thus, the vectors $w_{1,1}$, $w_{2,1}$, $w_{2,l}''$, $l \geq 2$, determine a column.

Case 2. Exactly one of the coordinates β_1^1 , β_1^2 , β_1^3 is distinct from a_1 .

Then we can assume that $\beta_1^1 \neq a_1$, $\beta_1^2 = \beta_1^3 = a_1$, as in the other case we would change the order of the appropriate coordinates. Take into account all vectors with codes $(1, l, 1, 1, 2)$, $l \geq 2$. They have the following form

$$w_{2,l}'' : a_1 \ \beta_1^1 \ \gamma_1^{2l-3} \ \gamma_1^{2l-2} \ a_2, \quad l \geq 2.$$

As $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\gamma_1^i = a_1$ for all $i \geq 1$. Thus, the vectors $w_{2,1}$, $w_{2,l}'$, $l \geq 2$, determine a column.

Case 3. Exactly two of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3$ are distinct from a_1 .

Then, by the same reason as before, we can assume that $\beta_1^1 \neq a_1$, $\beta_1^2 \neq a_1$, and $\beta_1^3 = a_1$. Take into account all vectors with codes $(1, l, 1, 1, 2)$ and $(1, 1, l, 1, 2)$, $l \geq 2$. As these vectors together with $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$ are distinguishable, they can be written in the following form

$$\begin{aligned} w_{2,l}' : & \quad a_1 \quad \beta_l^1 \quad \gamma_1^{2l-3} \quad \gamma_1^{2l-2} \quad a_2, \\ w_{3,l-1}' : & \quad a_1 \quad \delta_1^{2l-3} \quad \beta_l^2 \quad \delta_1^{2l-2} \quad a_2, \quad l \geq 2. \end{aligned}$$

Since the vectors $w_{2,l}'$, $l \geq 2$, and $w_{3,k}'$, $k \geq 1$, must be distinguishable, we have $\delta_1^i = \beta_1^1$ for $i = 1, 3, 5, \dots$ or $\gamma_1^i = \beta_1^2$ for $i = 1, 3, 5, \dots$. If the first possibility happens, then, as $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\delta_1^i = a_1$ for $i = 2, 4, \dots$. Thus, the vectors $w_{2,1}$, $w_{3,k}'$, $k \geq 1$, determine a column. If the second possibility takes place, then we have $\gamma_1^i = a_1$ for $i = 2, 4, \dots$, and the vectors $w_{2,1}$, $w_{2,k}'$, $k \geq 2$, determine a column.

Case 4. All of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3$ are distinct from a_1 .

Then consider all vectors belonging to T with codes $(1, l, 1, 1, 2)$, $l \geq 2$. Since each of these vectors is distinguishable from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they have the form

$$w_{2,l} : \quad a_1 \quad \beta_l^1 \quad \gamma_1^{2l-3} \quad \gamma_1^{2l-2} \quad a_2, \quad l \geq 2.$$

If $\gamma_1^i = \beta_1^2$ for $i = 1, 3, \dots$ and $\gamma_1^i = \beta_1^3$ for $i = 2, 4, \dots$, then the vectors $w_{2,l}$, $l \geq 1$, determine a column. Suppose that at least one of the above equalities does not happen. We can assume that $\gamma_1^1 \neq \beta_1^2$, as if $\gamma_1^1 = \beta_1^2$ and for example $\gamma_1^4 \neq \beta_1^3$, then we would change the order of the third and forth coordinates, and replace the code ε with $\varepsilon' := (\varepsilon_1, \tau \circ \varepsilon_2, \varepsilon_4, \varepsilon_3, \varepsilon_5)$, where τ is the transposition (23) . Take all vectors with codes $(1, 1, l, 1, 2)$, $l \geq 2$. They can be written in the form

$$w_{3,l-1} : \quad a_1 \quad \delta_1^{2l-3} \quad \beta_l^2 \quad \delta_1^{2l-2} \quad a_2, \quad l \geq 2.$$

The distinguishability of the vectors $w_{2,2}$ and $w_{3,k}$, $k \geq 1$ implies $\delta_1^i = \beta_1^1$ for $i = 1, 3, \dots$, as $\gamma_1^1 \neq \beta_1^2$. If now $\delta_1^i = \beta_1^3$ for $i = 2, 4, \dots$, then the vectors $w_{2,1}$, $w_{3,l}$, $l \geq 1$, determine a column. Therefore, we can assume that at least one of the coordinates δ_1^i , $i = 2, 4, \dots$, is distinct from β_1^3 . We can also assume

that $\delta_1^2 \neq \beta_1^3$, as if $\delta_1^2 = \beta_1^3$ and for example $\delta_1^4 \neq \beta_1^3$, then we would change the code replacing ε_3 by the composite $\tau \circ \varepsilon_3$, where $\tau = (23)$. Now, let us take into account all vectors with codes $(1, 1, 1, l, 2)$, $l \geq 2$. Since each of them must be distinguishable from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they can be written as follows

$$w_{4,l-1} : \quad a_1 \quad \varepsilon_1^{2l-3} \quad \varepsilon_1^{2l-2} \quad \beta_l^3 \quad a_2, \quad l \geq 2.$$

Since $\delta_1^2 \neq \beta_1^3$, by the distinguishability of the vectors $w_{3,1}$ and $w_{4,l}$, $l \geq 1$, we have $\varepsilon_1^i = \beta_1^2$ for $i = 2, 4, \dots$. Again, as the vectors $w_{4,l}$, $l \geq 1$, and $w_{2,2}$ are distinguishable, $\varepsilon_1^i = \beta_1^1$ for $i = 1, 3, \dots$ or $\gamma_1^2 = \beta_1^3$. If the first possibility takes place, then the vectors $w_{2,1}$, $w_{4,l}$, $l \geq 1$, determine a column. Hence, the second possibility has to be considered. Consequently, we obtain

$$w_{1,1} : \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1,$$

$$w_{1,2} : \quad a_2 \quad a_1 \quad a_1 \quad a_1 \quad \alpha_1^1,$$

$$w_{2,1} : \quad a_1 \quad \beta_1^1 \quad \beta_1^2 \quad \beta_1^3 \quad a_2,$$

$$w_{2,2} : \quad a_1 \quad \beta_2^1 \quad \gamma_1^1 \quad \beta_1^3 \quad a_2,$$

$$w_{3,1} : \quad a_1 \quad \beta_1^1 \quad \beta_2^2 \quad \delta_1^2 \quad a_2,$$

$$w_{4,1} : \quad a_1 \quad \varepsilon_1^1 \quad \beta_1^2 \quad \beta_2^3 \quad a_2.$$

Take into account all vectors with codes $(1, 1, 2, l, 2)$, $(1, l, 1, 2, 2)$, $(1, 2, l, 1, 2)$, $l \geq 2$. They have the form

$$w_{5,l-1} : \quad \eta_1^{2l-3} \quad \beta_1^1 \quad \beta_2^2 \quad \delta_l^2 \quad \eta_2^{2l-2},$$

$$w_{6,l-1} : \quad \mu_1^{2l-3} \quad \varepsilon_l^1 \quad \beta_1^2 \quad \beta_2^3 \quad \mu_2^{2l-2},$$

$$w_{7,l-1} : \quad \nu_1^{2l-3} \quad \beta_2^1 \quad \gamma_l^1 \quad \beta_1^3 \quad \nu_2^{2l-2}, \quad l \geq 2.$$

If $\eta_1^i = a_1$ for $i = 1, 3, \dots$ and $\eta_2^i = a_2$ for $i = 2, 4, \dots$, then the vectors $w_{3,1}$, $w_{5,l}$, $l \geq 1$, determine a column. If $\mu_1^i = a_1$ for $i = 1, 3, \dots$ and $\mu_2^i = a_2$ for $i = 2, 4, \dots$, then the vectors $w_{4,1}$, $w_{6,l}$, $l \geq 1$, determine a column. If $\nu_1^i = a_1$ for $i = 1, 3, \dots$ and $\nu_2^i = a_2$ for $i = 2, 4, \dots$, then the vectors $w_{2,2}$, $w_{7,l}$, $l \geq 1$, determine a column. Therefore, we can assume that each of the above three statements is false. Then, by the distinguishability of the vectors $w_{5,l}$, $w_{6,l}$, $w_{7,l}$, $l \geq 1$, and $w_{1,1}$, $w_{1,2}$, we have $\delta_1^2 = \varepsilon_1^1 = \gamma_1^1 = a_1$. In consequence, we

obtain

$$w_{2,1} : a_1 \ \beta_1^1 \ \beta_1^2 \ \beta_1^3 \ a_2,$$

$$w_{2,2} : a_1 \ \beta_2^1 \ a_1 \ \beta_1^3 \ a_2,$$

$$w_{3,1} : a_1 \ \beta_1^1 \ \beta_2^2 \ a_1 \ a_2,$$

$$w_{4,1} : a_1 \ a_1 \ \beta_1^2 \ \beta_2^3 \ a_2,$$

$$w_{5,1} : \eta_1^1 \ \beta_1^1 \ \beta_2^2 \ a_2 \ \eta_2^2.$$

Assume that $\eta_1^1 \neq a_1$, as if $\eta_1^1 = a_1$ and for example $\eta_2^2 \neq a_2$, then we would change the order of the first and fifth coordinates, and replace the code ε by $\varepsilon' := (\tau \circ \varepsilon_5, \varepsilon_2, \varepsilon_3, \varepsilon_4, \tau \circ \varepsilon_1)$, where $\tau = (1\ 2)$. Consider in addition all vectors with codes $(l, 1, 2, 2, 2)$, $l \geq 2$. They can be written in the form

$$w_{8,l-1} : \eta_l^1 \ \beta_1^1 \ \beta_2^2 \ a_2 \ \rho_2^{l-1}, \quad l \geq 2.$$

If $\rho_2^i = \eta_2^2$ for all $i \geq 1$, then the vectors $w_{5,1}$ and $w_{8,l}$, $l \geq 1$, determine a column. Therefore, we can assume that $\rho_2^1 \neq \eta_2^2$, as if $\rho_2^1 = \eta_2^2$ and for example $\rho_2^2 \neq \eta_2^2$, then we would change the code replacing ε_1 by ε'_1 , defined so that ε'_1 restricted to $\mathbb{R} \setminus (\eta_1^1 + \mathbb{Z})$ coincides with ε_1 , while ε'_1 restricted to $\eta_1^1 + \mathbb{Z}$ coincides with the composite $(2\ 3) \circ \varepsilon_1$. Moreover, we can also assume that $\eta_2^2 \neq a_2$, as if $\eta_2^2 = a_2$ and $\rho_2^1 \neq a_2$, then we would change the code similarly as above replacing ε_1 appropriately. Now, let us consider all vectors with codes $(1, 1, 2, 2, 1)$, $(1, 1, 2, 2, l)$, $l \geq 3$. They have the form

$$w_{9,1} : \sigma_1^1 \ \beta_1^1 \ \beta_2^2 \ a_2 \ \eta_1^2,$$

$$w_{9,l-1} : \sigma_1^{l-1} \ \beta_1^1 \ \beta_2^2 \ a_2 \ \eta_l^2, \quad l \geq 3.$$

Since $\rho_2^1 \neq \eta_2^2$, by the distinguishability of $w_{8,1}$ and $w_{9,l}$, $l \geq 1$, we have $\sigma_1^i = \eta_1^1$ for all $i \geq 1$. Thus, the vectors $w_{5,1}$ and $w_{9,l}$, $l \geq 1$, determine a column. $\square$

Lemma 1 *If a set T determining a cube tiling of $\mathbb{R}^6$ contains vectors of the form*

$$w_{1,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{1,2} : a_2 \ b_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{1,3} : a_1 \ a_2 \ b_1 \ a_1 \ a_1 \ a_1,$$

$$w_{1,4} : b_1 \ a_1 \ a_2 \ a_1 \ a_1 \ a_1,$$

then it contains vectors which determine a column.

Proof. Consider all vectors with codes $(2, l, 1, 1, 1, 1)$, $l \geq 2$. As these vectors together with $w_{1,1}$, $w_{1,2}$, $w_{1,3}$ and $w_{1,4}$ are distinguishable, they can be written in the following form

$$w_{2,l-1} : \quad a_2 \quad b_l \quad a_1 \quad \delta_1^{3l-5} \quad \delta_1^{3l-4} \quad \delta_1^{3l-3}, \quad l \geq 2.$$

If $\delta_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,2}$, $w_{2,l}$, $l \geq 1$, determine a column. Therefore, let us suppose that at least one of the coordinates δ_1^i , $i \geq 1$, is distinct from a_1 . We can assume that it is δ_1^1 , as if $\delta_1^1 = a_1$ and for example $\delta_1^5 \neq a_1$, then we would change the order of the fourth and fifth coordinates, and the natural code replacing ε_2 by ε'_2 defined so that ε'_2 restricted to $\mathbb{R} \setminus (b_1 + \mathbb{Z})$ coincides with ε_2 while ε'_2 restricted to $b_1 + \mathbb{Z}$ coincides with $(2\ 3) \circ \varepsilon_2$. Now, consider all vectors with codes $(2, 2, 1, l, 1, 1)$, $l \geq 2$. They have the form

$$w_{3,l-1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_1^1 \quad \varepsilon_1^{2l-3} \quad \varepsilon_1^{2l-2}, \quad l \geq 2.$$

If $\varepsilon_1^i = \delta_1^2$ for $i = 1, 3, \dots$ and $\varepsilon_1^i = \delta_1^3$ for $i = 2, 4, \dots$, then the vectors $w_{2,1}$ and $w_{3,l}$, $l \geq 1$, determine a column. Therefore, let us assume that at least one of the above equalities does not happen. We can assume that $\varepsilon_1^1 \neq \delta_1^2$, as if $\varepsilon_1^1 = \delta_1^2$ and for example $\varepsilon_1^4 \neq \delta_1^3$, then we would change the order of the fifth and sixth coordinates, and the natural code replacing ε_4 by ε'_4 so that $\varepsilon'_4 = (2\ 3) \circ \varepsilon_4$ on $\delta_1^1 + \mathbb{Z}$ and $\varepsilon'_4 = \varepsilon_4$ on the complement of $\delta_1^1 + \mathbb{Z}$. Moreover, we can also assume that $\delta_1^2 \neq a_1$. (If $\delta_1^2 = a_1$ and $\varepsilon_1^1 \neq a_1$, then we would change the natural code replacing ε_4 by ε'_4 so that $\varepsilon'_4 = (1\ 2) \circ \varepsilon_4$ on $\delta_1^1 + \mathbb{Z}$ and $\varepsilon'_4 = \varepsilon_4$ on the complement of $\delta_1^1 + \mathbb{Z}$.) Now, let us take all vectors with codes $(2, 2, 1, 1, l, 1)$, $l \geq 2$. They can be written in the form

$$w_{4,l-1} : \quad a_2 \quad b_2 \quad a_1 \quad \varphi_1^{2l-3} \quad \delta_1^2 \quad \varphi_1^{2l-2}, \quad l \geq 2.$$

By the distinguishability of the vectors $w_{3,1}$ and $w_{4,l}$, $l \geq 1$, we have $\varphi_1^i = \delta_1^1$ for $i = 1, 3, \dots$, as $\varepsilon_1^1 \neq \delta_1^2$. If now $\varphi_1^i = \delta_1^3$ for $i = 2, 4, \dots$, then $w_{2,1}$, $w_{4,l}$, $l \geq 1$, determine a column. Therefore, let us suppose that at least one of the coordinates φ_1^i , $i = 2, 4, \dots$, is distinct from δ_1^3 . We can assume that this is φ_1^2 . Moreover, we can also assume that $\delta_1^3 \neq a_1$. (If $\delta_1^3 = a_1$, and $\varphi_1^2 \neq a_1$, then we would replace ε_5 by ε'_5 so that $\varepsilon'_5 = (1\ 2) \circ \varepsilon_5$ on $\delta_1^1 + \mathbb{Z}$ and $\varepsilon'_5 = \varepsilon_5$ on the complement of the set $\delta_1^1 + \mathbb{Z}$.) Consider all vectors with codes $(2, 2, 1, 1, 1, l)$, $l \geq 2$. By the distinguishability, they have the form

$$w_{5,l-1} : \quad a_2 \quad b_2 \quad a_1 \quad \mu_1^{2l-3} \quad \mu_1^{2l-2} \quad \delta_1^3, \quad l \geq 2.$$

Since $\varphi_1^2 \neq \delta_1^3$, by the distinguishability of $w_{4,1}$ and $w_{5,l}$, $l \geq 1$, we have $\mu_1^i = \delta_1^2$ for $i = 2, 4, \dots$. Since the vectors $w_{3,1}$ and $w_{5,l}$, $l \geq 1$, must be distinguishable, we have $\mu_1^i = \delta_1^1$ for $i = 1, 3, \dots$ or $\varepsilon_1^2 = \delta_1^3$. If the first case takes place, then the vectors $w_{2,1}$, $w_{5,l}$, $l \geq 1$, determine a column. Hence, the second case must be considered. Then we obtain

$$w_{2,1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_1^1 \quad \delta_1^2 \quad \delta_1^3,$$

$$w_{3,1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_2^1 \quad \varepsilon_1^1 \quad \delta_1^3,$$

$$w_{4,1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_1^1 \quad \delta_2^2 \quad \varphi_1^2,$$

$$w_{5,1} : \quad a_2 \quad b_2 \quad a_1 \quad \mu_1^1 \quad \delta_1^2 \quad \delta_2^3.$$

Now, let us take into account all vectors with codes $(2, 2, 1, 2, l, 1)$, $l \geq 2$. By their distinguishability from $w_{2,1}$, $w_{3,1}$, $w_{4,1}$ and $w_{5,1}$, they have the following form

$$w_{6,l-1} : \quad \eta_2^{3l-5} \quad \eta_2^{3l-4} \quad \eta_1^{3l-3} \quad \delta_2^1 \quad \varepsilon_l^1 \quad \delta_1^3, \quad l \geq 2.$$

If $\eta_2^i = a_2$ for $i = 1, 4, \dots$, $\eta_2^i = b_2$ for $i = 2, 5, \dots$ and $\eta_1^i = a_1$ for $i = 3, 6, \dots$, then the vectors $w_{3,1}$, $w_{6,l}$, $l \geq 1$, determine a column. Therefore, let us assume that at least one of the above equalities does not happen. Then, by the distinguishability of $w_{6,l}$, $l \geq 1$, from $w_{1,1}$, $w_{1,2}$, $w_{1,3}$, $w_{1,4}$ we obtain $\varepsilon_1^1 = a_1$. Similarly, taking the vectors with codes $(2, 2, 1, 1, 2, l)$ and $(2, 2, 1, l, 1, 2)$, $l \geq 2$, we can show $\varphi_1^2 = a_1$ and $\mu_1^1 = a_1$. As a result, we have

$$w_{3,1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_2^1 \quad a_1 \quad \delta_1^3,$$

$$w_{4,1} : \quad a_2 \quad b_2 \quad a_1 \quad \delta_1^1 \quad \delta_2^2 \quad a_1,$$

$$w_{5,1} : \quad a_2 \quad b_2 \quad a_1 \quad a_1 \quad \delta_1^2 \quad \delta_2^3.$$

If we repeat the above reasoning starting with vectors with codes $(1, 2, l, 1, 1, 1)$, $l \geq 2$, instead of $(2, l, 1, 1, 1, 1)$, then we can add up the following vectors

$$w_{7,1} : \quad a_1 \quad a_2 \quad b_2 \quad \lambda_2^1 \quad a_1 \quad \lambda_1^3,$$

$$w_{8,1} : \quad a_1 \quad a_2 \quad b_2 \quad \lambda_1^1 \quad \lambda_2^2 \quad a_1,$$

$$w_{9,1} : \quad a_1 \quad a_2 \quad b_2 \quad a_1 \quad \lambda_1^2 \quad \lambda_2^3.$$

However, it can happen that in order to get such a system the change of the order of fourth, fifth and sixth coordinates as well as the change of the code

are to be performed. Such changes can affect the vectors $w_{3,1}$, $w_{4,1}$, $w_{5,1}$ but not in a substantial way. (The scheme remains unchanged.)

Now, we repeat the whole procedure once again beginning from the vectors with codes $(l, 1, 2, 1, 1, 1)$, $l \geq 2$. Then we can add up the following vectors

$$w_{10,1} : b_2 \ a_1 \ a_2 \ \nu_2^1 \ a_1 \ \nu_1^3,$$

$$w_{11,1} : b_2 \ a_1 \ a_2 \ \nu_1^1 \ \nu_2^2 \ a_1,$$

$$w_{12,1} : b_2 \ a_1 \ a_2 \ a_1 \ \nu_1^2 \ \nu_2^3.$$

As previously, the vectors $w_{3,1}$, $w_{4,1}$, $w_{5,1}$ and $w_{7,1}$, $w_{8,1}$, $w_{9,1}$ can be affected. The resulting system can be written as follows

$$w_{3,1} : a_2 \ b_2 \ a_1 \ A_2^1 \ a_1 \ A_1^3,$$

$$w_{4,1} : a_2 \ b_2 \ a_1 \ A_1^1 \ A_2^2 \ a_1,$$

$$w_{5,1} : a_2 \ b_2 \ a_1 \ a_1 \ A_1^2 \ A_2^3,$$

$$w_{7,1} : a_1 \ a_2 \ b_2 \ B_2^1 \ a_1 \ B_1^3,$$

$$w_{8,1} : a_1 \ a_2 \ b_2 \ B_1^1 \ B_2^2 \ a_1,$$

$$w_{9,1} : a_1 \ a_2 \ b_2 \ a_1 \ B_1^2 \ B_2^3,$$

$$w_{10,1} : b_2 \ a_1 \ a_2 \ \nu_2^1 \ a_1 \ \nu_1^3,$$

$$w_{11,1} : b_2 \ a_1 \ a_2 \ \nu_1^1 \ \nu_2^2 \ a_1,$$

$$w_{12,1} : b_2 \ a_1 \ a_2 \ a_1 \ \nu_1^2 \ \nu_2^3,$$

where capital Roman letters with lower and upper indices have a similar meaning to that of Greek letters except that the lower index may not be equal to the code value. Precisely, they are different from all a_i , $i \geq 1$; any two symbols that differ only by lower indices represent reals whose distance is a non-zero integer; e.g. $A_1^1 - A_2^1 \in \mathbb{Z} \setminus \{0\}$.

Now, let us consider all vectors with codes $(2, 1, 1, l, 1, 1)$, $l \geq 2$. Since $w_{1,2}$ has its code equal to $(2, 1, 1, 1, 1, 1)$, these vectors are as follows

$$w_{13,l-1} : \rho_2^{5l-9} \ \rho_1^{5l-8} \ \rho_1^{5l-7} \ a_l \ \rho_1^{5l-6} \ \rho_1^{5l-5}, \quad l \geq 2.$$

Similarly, the vectors with codes $(2, 1, 1, 1, l, 1)$, $l \geq 2$, and $(2, 1, 1, 1, 1, l)$, $l \geq 2$, can be written as follows

$$\begin{aligned} w_{14, l-1} : & \sigma_2^{5l-9} \quad \sigma_1^{5l-8} \quad \sigma_1^{5l-7} \quad \sigma_1^{5l-6} \quad a_l \quad \sigma_1^{5l-5}, \\ w_{15, l-1} : & \tau_2^{5l-9} \quad \tau_1^{5l-8} \quad \tau_1^{5l-7} \quad \tau_1^{5l-6} \quad \tau_1^{5l-5} \quad a_l, \end{aligned} \quad l \geq 2.$$

If $\rho_1^i = a_1$ for $i = 4, 9, \dots$ and $i = 5, 10, \dots$, then by the distinguishability of $w_{13, l}$, $l \geq 1$, from $w_{10, 1}$, $w_{3, 1}$ and $w_{7, 1}$, we have $\rho_2^i = a_2$ for $i = 1, 6, \dots$, $\rho_1^i = b_1$ for $i = 2, 7, \dots$ and $\rho_1^i = a_1$ for $i = 3, 8, \dots$. Then $w_{1, 2}$, $w_{13, l}$, $l \geq 1$, determine a column. If $\sigma_1^i = a_1$ for $i = 4, 9, \dots$ and $i = 5, 10, \dots$, or $\tau_1^i = a_1$ for $i = 4, 9, \dots$ and $i = 5, 10, \dots$, then, in the same way, we show that the set T contains vectors determining a column. Suppose that none of these situations happen. We can assume that $\rho_1^4 \neq a_1$. (If for example $\rho_1^{10} \neq a_1$ and $\rho_1^4 = a_1$, then we would change the order of the fifth and sixth coordinates, and replace ε_4 by ε'_4 so that $\varepsilon'_4 = (2\ 3) \circ \varepsilon_4$ on $a_1 + \mathbb{Z}$ and $\varepsilon'_4 = \varepsilon_4$ on the complement of $a_1 + \mathbb{Z}$.) Then by the distinguishability of $w_{13, 1}$ and $w_{14, l}$, $l \geq 1$, we have $\sigma_1^i = a_1$ for $i = 4, 9, \dots$. We can assume that $\sigma_1^5 \neq a_1$, as $\sigma_1^i \neq a_1$ for some $i \in \{5, 10, \dots\}$. By the distinguishability of $w_{14, 1}$ and $w_{15, l}$, $l \geq 1$, we have $\tau_1^i = a_1$ for $i = 5, 10, \dots$. Then we can assume that $\tau_1^4 \neq a_1$. By the distinguishability of $w_{15, 1}$ and $w_{13, l}$, $l \geq 1$, we have $\rho_1^i = a_1$ for $i = 5, 10, \dots$. As a result, we obtain

$$\begin{aligned} w_{13, 1} : & \rho_2^1 \quad \rho_1^2 \quad \rho_1^3 \quad a_2 \quad \rho_1^4 \quad a_1, \\ w_{14, 1} : & \sigma_2^1 \quad \sigma_1^2 \quad \sigma_1^3 \quad a_1 \quad a_2 \quad \sigma_1^5, \\ w_{15, 1} : & \tau_2^1 \quad \tau_1^2 \quad \tau_1^3 \quad \tau_1^4 \quad a_1 \quad a_2. \end{aligned}$$

Now, let us take into account all vectors with codes $(2, 1, 1, 2, l, 1)$, $l \geq 2$. By their distinguishability from $w_{13, 1}$, $w_{14, 1}$ and $w_{15, 1}$, they have the following form

$$w_{16, l-1} : \theta_2^{3l-5} \quad \theta_1^{3l-4} \quad \theta_1^{3l-3} \quad a_2 \quad \rho_l^4 \quad a_1, \quad l \geq 2.$$

By the distinguishability of $w_{13, 1}$ and $w_{16, l}$, $l \geq 1$, from $w_{10, 1}$, $w_{3, 1}$ and $w_{7, 1}$, we have $\rho_1^3 = a_1$, $\rho_1^2 = b_1$, $\rho_2^1 = a_2$, $\theta_1^i = a_1$ for $i = 3, 6, \dots$, $\theta_1^i = b_1$ for $i = 2, 5, \dots$ and $\theta_2^i = a_2$ for $i = 1, 4, \dots$. Thus, the vectors $w_{13, 1}$ and $w_{16, l}$, $l \geq 1$, determine a column. $\square$

Theorem 8 *Every cube tiling of $\mathbb{R}^6$ contains a column.*

Proof. Let T be an arbitrary set which determines a cube tiling of $\mathbb{R}^6$. By Theorems 3 and 7, the set T contains vectors which determine a 4-column.

Passing to an isomorphic system if necessary, we can assume that the vectors determining our 4-column are as follows

$$w_{1,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{1,l} : a_l \ a_1 \ a_1 \ a_1 \ a_1 \ \alpha_1^{l-1}, \quad l \geq 2.$$

If $\alpha_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{1,k}$, $k \geq 1$, determine a column. Suppose that $\alpha_1^i \neq a_1$ for some $i \geq 1$. We can assume that $\alpha_1^1 \neq a_1$. Consider all vectors with codes $(1, *, *, *, *, l)$, $l \geq 2$, where $*$ can take any value from $\mathbb{N}$, such that their middle coordinates (second, third, fourth and fifth) are different from a_l , $l \geq 2$. Let us pick a vector from among them whose middle coordinates differ from a_1 at as many places as possible. Similarly as in the proof of Theorem 6 we can assume that this vector has the code $(1, 1, 1, 1, 1, 2)$. By its distinguishability from $w_{1,1}$ and $w_{1,2}$, it can be written in the form

$$w_{2,1} : a_1 \ \beta_1^1 \ \beta_1^2 \ \beta_1^3 \ \beta_1^4 \ a_2.$$

Five cases have to be considered:

Case 1. $\beta_1^1 = \beta_1^2 = \beta_1^3 = \beta_1^4 = a_1$.

Then take all vectors with codes $(1, 1, 1, 1, 1, l)$, $l \geq 3$. Since each of them must be distinguishable from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they have the form

$$w_{2,l-1}''' : a_1 \ \beta_1^{4l-7} \ \beta_1^{4l-6} \ \beta_1^{4l-5} \ \beta_1^{4l-4} \ a_l, \quad l \geq 3.$$

Since $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\beta_1^i = a_1$ for all $i \geq 5$. Thus, the vectors $w_{2,1}$, $w_{2,l}'''$, $l \geq 2$, determine a column.

Case 2. Exactly three of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3, \beta_1^4$ are equal to a_1 .

Then we can assume that $\beta_1^2 = \beta_1^3 = \beta_1^4 = a_1$, $\beta_1^1 \neq a_1$, as in the other case we would change the order of the appropriate coordinates. (If $\beta_1^1 = a_1$ and for example $\beta_1^4 \neq a_1$, then we exchange the second coordinate with the fifth.) Take into account all vectors with codes $(1, l, 1, 1, 1, 2)$, $l \geq 2$. By their distinguishability from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they have the following form

$$w_{2,l}''' : a_1 \ \beta_1^1 \ \gamma_1^{3l-5} \ \gamma_1^{3l-4} \ \gamma_1^{3l-3} \ a_2, \quad l \geq 2.$$

As $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\gamma_1^i = a_1$ for all $i \geq 1$. Thus, the vectors $w_{2,1}$, $w_{2,l}'''$, $l \geq 2$, determine a column.

Case 3. Exactly two of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3, \beta_1^4$ are equal to a_1 .

Then, by the same reasoning as before, we can assume that $\beta_1^3 = \beta_1^4 = a_1$, $\beta_1^1 \neq a_1$ and $\beta_1^2 \neq a_1$. Consider all vectors with codes $(1, l, 1, 1, 1, 2)$ and $(1, 1, l, 1, 1, 2)$, $l \geq 2$. They can be written in the form

$$\begin{aligned} w''_{2,l} : & \quad a_1 \quad \beta_l^1 \quad \gamma_1^{3l-5} \quad \gamma_1^{3l-4} \quad \gamma_1^{3l-3} \quad a_2, \\ w''_{3,l-1} : & \quad a_1 \quad \delta_1^{3l-5} \quad \beta_l^2 \quad \delta_1^{3l-4} \quad \delta_1^{3l-3} \quad a_2, \quad l \geq 2. \end{aligned}$$

Since the vectors $w''_{2,k}$, $k \geq 2$, and $w''_{3,l}$, $l \geq 1$, must be distinguishable, we have $\delta_1^i = \beta_1^1$ for $i = 1, 4, \dots$, or $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$. If the first case takes place, we have $\delta_1^i = a_1$ for $i = 2, 5, \dots$ and $i = 3, 6, \dots$, as $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 . Then the vectors $w_{2,1}$, $w''_{3,l}$, $l \geq 1$, determine a column. Otherwise, $\gamma_1^i = a_1$ for $i = 2, 5, \dots$ and $i = 3, 6, \dots$, and the vectors $w_{2,1}$, $w''_{2,k}$, $k \geq 2$, determine a column.

Case 4. Exactly one of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3, \beta_1^4$ is equal to a_1 .

Then we can suppose that $\beta_1^4 = a_1$, $\beta_1^1 \neq a_1$, $\beta_1^2 \neq a_1$ and $\beta_1^3 \neq a_1$. Consider all vectors with codes $(1, l, 1, 1, 1, 2)$, $l \geq 2$. By their distinguishability from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they can be written as follows

$$w'_{2,l} : \quad a_1 \quad \beta_l^1 \quad \gamma_1^{3l-5} \quad \gamma_1^{3l-4} \quad \gamma_1^{3l-3} \quad a_2, \quad l \geq 2.$$

If $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$ and $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$, then we have $\gamma_1^i = a_1$ for $i = 3, 6, \dots$, as $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 . Thus, the vectors $w_{2,1}$, $w'_{2,l}$, $l \geq 2$, determine a column. Therefore, let us suppose that at least one of the above equalities does not happen. We can assume that $\gamma_1^1 \neq \beta_1^2$. (If $\gamma_1^1 = \beta_1^2$ and for example $\gamma_1^5 \neq \beta_1^3$, then we would change the order of the third and fourth coordinates, and the code replacing ε_2 by ε'_2 so that $\varepsilon'_2 = (23) \circ \varepsilon_2$ on $\beta_1^1 + \mathbb{Z}$ and $\varepsilon'_2 = \varepsilon_2$ on the complement of $\beta_1^1 + \mathbb{Z}$.) Take into account all vectors with codes $(1, 1, l, 1, 1, 2)$, $l \geq 2$. They have the form

$$w'_{3,l-1} : \quad a_1 \quad \delta_1^{3l-5} \quad \beta_l^2 \quad \delta_1^{3l-4} \quad \delta_1^{3l-3} \quad a_2, \quad l \geq 2.$$

Since $\gamma_1^1 \neq \beta_1^2$ and the vectors $w'_{2,2}$ and $w'_{3,l}$, $l \geq 1$, are distinguishable, we have $\delta_1^i = \beta_1^1$ for $i = 1, 4, \dots$. If now $\delta_1^i = \beta_1^3$ for $i = 2, 5, \dots$, then, since $\beta_1^4 = a_1$, and $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\delta_1^i = a_1$ for $i = 3, 6, \dots$. Thus, the vectors $w_{2,1}$, $w'_{3,l}$, $l \geq 1$, determine a column. Therefore, let us assume that $\delta_1^2 \neq \beta_1^3$. (If $\delta_1^2 = \beta_1^3$, and for example $\delta_1^4 \neq \beta_1^1$, then we would replace the code ε_3 by ε'_3 so that $\varepsilon'_3 = (23) \circ \varepsilon_3$ on $\beta_1^2 + \mathbb{Z}$ and $\varepsilon'_3 = \varepsilon_3$ on the complement of $\beta_1^2 + \mathbb{Z}$.)

Take all vectors with codes $(1, 1, 1, l, 1, 2)$, $l \geq 2$. They can be written in the form

$$w'_{4, l-1} : a_1 \quad \varepsilon_1^{3l-5} \quad \varepsilon_1^{3l-4} \quad \beta_l^3 \quad \varepsilon_1^{3l-3} \quad a_2, \quad l \geq 2.$$

Since $\delta_1^2 \neq \beta_1^3$, by the distinguishability of the vectors $w'_{3,1}$ and $w'_{4,l}$, $l \geq 1$, we have $\varepsilon_1^i = \beta_1^2$ for $i = 2, 5, \dots$. Since $w'_{2,2}$ and $w'_{4,l}$, $l \geq 1$, are distinguishable, we have $\varepsilon_1^i = \beta_1^1$ for $i = 1, 4, \dots$ or $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$. If the first possibility happens, then $\varepsilon_1^i = a_1$ for $i = 3, 6, \dots$, and the vectors $w_{2,1}$, $w'_{4,l}$, $l \geq 1$, determine a column. Hence, $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$. Consequently, we obtain

$$w_{1,1} : a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1,$$

$$w_{1,2} : a_2 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad \alpha_1^1,$$

$$w_{2,1} : a_1 \quad \beta_1^1 \quad \beta_1^2 \quad \beta_1^3 \quad a_1 \quad a_2,$$

$$w'_{2,2} : a_1 \quad \beta_2^1 \quad \gamma_1^1 \quad \beta_1^3 \quad \gamma_1^3 \quad a_2,$$

$$w'_{3,1} : a_1 \quad \beta_1^1 \quad \beta_2^2 \quad \delta_1^2 \quad \delta_1^3 \quad a_2,$$

$$w'_{4,1} : a_1 \quad \varepsilon_1^1 \quad \beta_1^2 \quad \beta_2^3 \quad \varepsilon_1^3 \quad a_2.$$

If now $\gamma_1^1 \neq a_1$, then take all vectors with codes $(1, 2, l, 1, 1, 2)$, $l \geq 2$. They have the form

$$w'_{5, l-1} : a_1 \quad \beta_2^1 \quad \gamma_l^1 \quad \beta_1^3 \quad \varphi_1^{l-1} \quad a_2, \quad l \geq 2.$$

Since $\gamma_1^1 \neq a_1$, $\beta_1^3 \neq a_1$, and $w_{2,1}$ has the smallest possible number of the middle coordinates equal to a_1 , we have $\gamma_1^3 = a_1$ and $\varphi_1^i = a_1$ for all $i \geq 1$. Thus, the vectors $w'_{2,2}$, $w'_{5,l}$, $l \geq 1$, determine a column. Similarly, we show that the set T contains the vectors determining a column, if $\delta_1^2 \neq a_1$ or $\varepsilon_1^1 \neq a_1$. Therefore, the possibility $\gamma_1^1 = \delta_1^2 = \varepsilon_1^1 = a_1$ must be considered. Then we have

$$w_{2,1} : a_1 \quad \beta_1^1 \quad \beta_1^2 \quad \beta_1^3 \quad a_1 \quad a_2,$$

$$w'_{2,2} : a_1 \quad \beta_2^1 \quad a_1 \quad \beta_1^3 \quad \gamma_1^3 \quad a_2,$$

$$w'_{3,1} : a_1 \quad \beta_1^1 \quad \beta_2^2 \quad a_1 \quad \delta_1^3 \quad a_2,$$

$$w'_{4,1} : a_1 \quad a_1 \quad \beta_1^2 \quad \beta_2^3 \quad \varepsilon_1^3 \quad a_2.$$

If now $\gamma_1^3 = \delta_1^3 = \varepsilon_1^3 = a_1$, then we rename β_1^1 , β_1^2 , β_1^3 replacing them by a_1

and vice versa. Then we obtain

$$w_{2,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1 \ a_2,$$

$$w'_{2,2} : a_1 \ a_2 \ \beta_1^2 \ a_1 \ a_1 \ a_2,$$

$$w'_{3,1} : a_1 \ a_1 \ a_2 \ \beta_1^3 \ a_1 \ a_2,$$

$$w'_{4,1} : a_1 \ \beta_1^1 \ a_1 \ a_2 \ a_1 \ a_2.$$

Subsequently, we change the order of the coordinates applying the cyclic permutation $(1\ 4\ 3\ 2)$, and change the code ε_6 by ε'_6 so that $\varepsilon'_6 = (1\ 2) \circ \varepsilon_6$ on $a_1 + \mathbb{Z}$ and $\varepsilon'_6 = \varepsilon_6$ on the complement of $a_1 + \mathbb{Z}$. As a result, we obtain

$$w_{2,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w'_{2,2} : a_2 \ \beta_1^2 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w'_{3,1} : a_1 \ a_2 \ \beta_1^3 \ a_1 \ a_1 \ a_1,$$

$$w'_{4,1} : \beta_1^1 \ a_1 \ a_2 \ a_1 \ a_1 \ a_1.$$

By Lemma 1, the set T contains vectors determining a column. Therefore, let us assume that $\gamma_1^3 \neq a_1$. Consider all vectors with codes $(1, 2, 1, 1, l, 2)$, $l \geq 2$. By the distinguishability, they are as follows:

$$w'_{6,l-1} : a_1 \ \beta_2^1 \ \eta_1^{2l-3} \ \eta_1^{2l-2} \ \gamma_l^3 \ a_2, \quad l \geq 2.$$

If $\gamma_1^3 \neq \varepsilon_1^3$, then by the distinguishability of the vectors $w'_{6,l}$, $l \geq 1$, and $w'_{4,1}$ we have $\eta_1^i = \beta_1^3$ for $i = 2, 4, \dots$. We also have $\eta_1^i = a_1$ for $i = 1, 3, \dots$, as w_2^1 has the smallest possible number of the middle coordinates equal to a_1 . Thus, the vectors $w'_{2,2}$, $w'_{6,l}$, $l \geq 1$, determine a column. If $\gamma_1^3 = \varepsilon_1^3$, then in particular $\varepsilon_1^3 \neq a_1$. Take all vectors with codes $(1, 1, 1, 2, l, 2)$, $l \geq 2$. By the distinguishability, they have the form

$$w'_{7,l-1} : a_1 \ \mu_1^{2l-3} \ \mu_1^{2l-2} \ \beta_2^3 \ \varepsilon_l^3 \ a_2, \quad l \geq 2.$$

If $\varepsilon_1^3 \neq \delta_1^3$, then by the distinguishability of the vectors $w'_{3,1}$ and $w'_{7,l}$, $l \geq 1$, we have $\mu_1^i = \beta_1^2$ for $i = 2, 4, \dots$. Then $\mu_1^i = a_1$ for $i = 1, 3, \dots$ and the vectors $w'_{4,1}$, $w'_{7,l}$, $l \geq 1$, determine a column. In the same way we show that the set T contains vectors determining a column, if we assume that $\delta_1^3 \neq a_1$ or $\varepsilon_1^3 \neq a_1$. Now, let us assume that $\gamma_1^3 = \delta_1^3 = \varepsilon_1^3 \neq a_1$. Let us take into

account vectors $w'_{2,2}$, $w'_{3,1}$ and $w'_{4,1}$. It should be clear that passing to an appropriate isomorphic system we can assume that

$$w'_{2,2} : \begin{matrix} a_1 & a_1 & a_1 & a_1 & a_1 & a_1 \end{matrix},$$

$$w'_{3,1} : \begin{matrix} a_1 & a_2 & \beta_2^2 & \beta_1^3 & a_1 & a_1 \end{matrix},$$

$$w'_{4,1} : \begin{matrix} a_1 & \beta_2^1 & \beta_1^2 & a_2 & a_1 & a_1 \end{matrix}.$$

Now, let us consider all vectors with codes $(1, 1, l, 1, 1, 1)$, $l \geq 2$. By their distinguishability from $w'_{2,2}$, $w'_{3,1}$ and $w'_{4,1}$, they can be written as follows:

$$w'_{8,l-1} : \begin{matrix} \nu_1^{3l-5} & a_1 & a_l & a_1 & \nu_1^{3l-4} & \nu_1^{3l-3} \end{matrix}, \quad l \geq 2.$$

If $\nu_1^i = a_1$ for all $i \geq 1$, then the vectors $w'_{2,2}$, $w'_{8,l}$, $l \geq 1$, determine a column. Therefore, let us suppose that at least one of the coordinates ν_1^i , $i \geq 1$, is distinct from a_1 . We can assume that $\nu_1^1 \neq a_1$. (If not, then we would change the order of coordinates and the code ε_3 appropriately.) Take into account all vectors with codes $(l, 1, 2, 1, 1, 1)$, $l \geq 2$. They have the form

$$w'_{9,l-1} : \begin{matrix} \nu_l^1 & a_1 & a_2 & a_1 & \rho_1^{2l-3} & \rho_1^{2l-2} \end{matrix}, \quad l \geq 2.$$

If $\rho_1^i = \nu_1^2$ for $i = 1, 3, \dots$ and $\rho_1^i = \nu_1^3$ for $i = 2, 4, \dots$, then the vectors $w'_{8,1}$, $w'_{9,l}$, $l \geq 1$, determine a column. Therefore, let us assume that $\rho_1^1 \neq \nu_1^2$. (If not, then we would change the order of the fifth and sixth coordinates and the code ε_1 appropriately.) Then we can also assume that $\nu_1^2 \neq a_1$. (If $\rho_1^1 \neq a_1$ and $\nu_1^2 = a_1$, then we would replace the code ε_1 by ε'_1 so that $\varepsilon'_1 = (1\ 2) \circ \varepsilon_1$ on $\nu_1^1 + \mathbb{Z}$ and $\varepsilon'_1 = \varepsilon_1$ on the complement of $\nu_1^1 + \mathbb{Z}$.) Consider all vectors with codes $(1, 1, 2, 1, l, 1)$, $l \geq 2$. They are as follows

$$w'_{10,l-1} : \begin{matrix} \sigma_1^{2l-3} & a_1 & a_2 & a_1 & \nu_l^2 & \sigma_1^{2l-2} \end{matrix}, \quad l \geq 2.$$

Since $\rho_1^1 \neq \nu_1^2$ and the vectors $w'_{9,1}$ and $w'_{10,l}$, $l \geq 1$, are distinguishable, we have $\sigma_1^i = \nu_1^1$ for $i = 1, 3, \dots$. If now $\sigma_1^i = \nu_1^3$ for $i = 2, 4, \dots$, then the vectors $w'_{8,1}$, $w'_{10,l}$, $l \geq 1$, determine a column. Therefore, let us assume that $\sigma_1^2 \neq \nu_1^3$. (If not, then we would change the code ε_5 appropriately.) Then we can also assume that $\nu_1^3 \neq a_1$. Take all vectors with codes $(1, 1, 2, 1, 1, l)$, $l \geq 2$. They can be written in the following form

$$w'_{11,l-1} : \begin{matrix} \tau_1^{2l-3} & a_1 & a_2 & a_1 & \tau_1^{2l-2} & \nu_l^3 \end{matrix}, \quad l \geq 2.$$

By the distinguishability of the vectors $w'_{11,l}$, $l \geq 1$, and $w'_{10,1}$ we have $\tau_1^i = \nu_1^2$ for $i = 2, 4, \dots$, as $\sigma_1^2 \neq \nu_1^3$. Since $w'_{9,1}$ and $w'_{11,l}$, $l \geq 1$, are distinguishable,

we have $\tau_1^i = \nu_1^1$ for $i = 1, 3, \dots$ or $\rho_1^i = \nu_1^3$ for $i = 2, 4, \dots$. If the first possibility happens, then the vectors $w'_{8,1}, w'_{11,l}, l \geq 1$, determine a column. Hence, it remains to consider the second possibility. Consequently, we obtain

$$w'_{8,1} : \quad \nu_1^1 \quad a_1 \quad a_2 \quad a_1 \quad \nu_1^2 \quad \nu_1^3,$$

$$w'_{9,1} : \quad \nu_2^1 \quad a_1 \quad a_2 \quad a_1 \quad \rho_1^1 \quad \nu_1^3,$$

$$w'_{10,1} : \quad \nu_1^1 \quad a_1 \quad a_2 \quad a_1 \quad \nu_2^2 \quad \sigma_1^2,$$

$$w'_{11,1} : \quad \tau_1^1 \quad a_1 \quad a_2 \quad a_1 \quad \nu_1^2 \quad \nu_2^3.$$

Passing to an appropriate isomorphic system, as has been done before, we can assume that the latter system of vectors has the form

$$w'_{8,1} : \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1,$$

$$w'_{9,1} : \quad a_2 \quad \rho_1^1 \quad a_1 \quad a_1 \quad a_1 \quad a_1,$$

$$w'_{10,1} : \quad a_1 \quad a_2 \quad \sigma_1^2 \quad a_1 \quad a_1 \quad a_1,$$

$$w'_{11,1} : \quad \tau_1^1 \quad a_1 \quad a_2 \quad a_1 \quad a_1 \quad a_1.$$

By Lemma 1, the set T contains vectors determining a column.

Case 5. All of the coordinates $\beta_1^1, \beta_1^2, \beta_1^3, \beta_1^4$ are distinct from a_1 .

Let us remind that

$$w_{1,1} : \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1,$$

$$w_{1,2} : \quad a_2 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad \alpha_1^1,$$

$$w_{2,1} : \quad a_1 \quad \beta_1^1 \quad \beta_1^2 \quad \beta_1^3 \quad \beta_1^4 \quad a_2.$$

Now, let us consider all vectors with codes $(1, l, 1, 1, 1, 2)$, $(1, 1, l, 1, 1, 2)$, $(1, 1, 1, l, 1, 2)$ and $(1, 1, 1, 1, l, 2)$, $l \geq 2$. By their distinguishability from $w_{1,1}$, $w_{1,2}$ and $w_{2,1}$, they have the form

$$w_{2,l} : \quad a_1 \quad \beta_l^1 \quad \gamma_1^{3l-5} \quad \gamma_1^{3l-4} \quad \gamma_1^{3l-3} \quad a_2,$$

$$w_{3,l-1} : \quad a_1 \quad \delta_1^{3l-5} \quad \beta_l^2 \quad \delta_1^{3l-4} \quad \delta_1^{3l-3} \quad a_2,$$

$$w_{4,l-1} : \quad a_1 \quad \varepsilon_1^{3l-5} \quad \varepsilon_1^{3l-4} \quad \beta_l^3 \quad \varepsilon_1^{3l-3} \quad a_2,$$

$$w_{5,l-1} : \quad a_1 \quad \eta_1^{3l-5} \quad \eta_1^{3l-4} \quad \eta_1^{3l-3} \quad \beta_l^4 \quad a_2, \quad l \geq 2.$$

We prove now that there is $\kappa \in \{\gamma, \delta, \varepsilon, \eta\}$ such that at least two of the families $\{\kappa_1^{3l-5} : l \geq 2\}$, $\{\kappa_1^{3l-4} : l \geq 2\}$, $\{\kappa_1^{3l-3} : l \geq 2\}$ are singletons $\{\beta_1^{i-1}\}$, where i relates to the i -th coordinate of the vectors under consideration. The distinguishability of the vectors $w_{2,l}$ and $w_{3,l-1}$, $l \geq 2$, implies $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$ or $\delta_1^i = \beta_1^1$ for $i = 1, 4, \dots$. We can assume that the first case takes place. (If $\gamma_1^i \neq \beta_1^2$ for some $i \in \{1, 4, \dots\}$, then $\delta_1^i = \beta_1^1$ for $i = 1, 4, \dots$ and we would change the order of the second and third coordinates.) By the distinguishability of the vectors $w_{2,l}$ and $w_{4,l-1}$, $l \geq 2$, we have $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$ or $\varepsilon_1^i = \beta_1^1$ for $i = 1, 4, \dots$. If the first possibility happens, then each vector $w_{2,l}$, $l \geq 2$, has the third coordinate equal to β_1^2 and the fourth equal to β_1^3 . Therefore, let us assume that $\varepsilon_1^i = \beta_1^1$ for $i = 1, 4, \dots$. Since $w_{3,l}$ and $w_{4,l}$, $l \geq 1$, are distinguishable, we have $\delta_1^i = \beta_1^3$ for $i = 2, 5, \dots$ or $\varepsilon_1^i = \beta_1^2$ for $i = 2, 5, \dots$. In the second case each vector $w_{4,l}$, $l \geq 1$, has its second and third coordinates equal to β_1^1 and β_1^2 , respectively. Hence $\delta_1^i = \beta_1^3$ for $i = 2, 5, \dots$. The distinguishability of $w_{3,l}$ and $w_{5,l}$, $l \geq 1$, implies $\eta_1^i = \beta_1^2$ for $i = 2, 5, \dots$ or $\delta_1^i = \beta_1^4$ for $i = 3, 6, \dots$. If the second possibility happens, then each vector $w_{3,l}$, $l \geq 1$, has its fourth and fifth coordinates equal to β_1^3 and β_1^4 , respectively. Therefore, we can assume that $\eta_1^i = \beta_1^2$ for $i = 2, 5, \dots$. Since $w_{2,l}$ and $w_{5,l-1}$, $l \geq 2$, are distinguishable, we have $\gamma_1^i = \beta_1^4$ for $i = 3, 6, \dots$ or $\eta_1^i = \beta_1^1$ for $i = 1, 4, \dots$. In the first case each vector of $w_{2,l}$, $l \geq 2$, has its third and fifth coordinates equal to β_1^2 and β_1^4 , respectively. In the second case each vector of $w_{5,l}$, $l \geq 1$, has the second coordinate equal to β_1^1 and the third equal to β_1^2 .

As a result, we can assume that for the block $w_{2,l}$, $l \geq 2$, at least two of the following equations hold:

$$\{\gamma_1^{3l-5} : l \geq 2\} = \{\beta_1^2\}, \quad \{\gamma_1^{3l-4} : l \geq 2\} = \{\beta_1^3\}, \quad \{\gamma_1^{3l-3} : l \geq 2\} = \{\beta_1^4\}.$$

(If it were for example the block $w_{4,l}$, $l \geq 1$ instead of $w_{2,l}$, $l \geq 2$, then we would change the order of the second and fourth coordinates.) We can also assume that $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$ and $\gamma_1^i = \beta_1^4$ for $i = 3, 6, \dots$. (If $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$ and $\gamma_1^i = \beta_1^3$ for $i = 2, 5, \dots$, then we would change the order of the third and fifth coordinates. If $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$ and $\gamma_1^i = \beta_1^4$ for $i = 3, 6, \dots$, then we would change the order of the third and fourth coordinates.) Then we can also assume that $\delta_1^i = \beta_1^4$ for $i = 3, 6, \dots$. (If $\delta_1^i \neq \beta_1^4$ for some $i \in \{3, 6, \dots\}$, but $\delta_1^i = \beta_1^3$ for $i = 2, 5, \dots$, then we would change the order of the fourth and fifth coordinates. If $\delta_1^i \neq \beta_1^4$ for some $i \in \{3, 6, \dots\}$ and $\delta_1^i \neq \beta_1^3$ for some $i \in \{2, 5, \dots\}$, then by the distinguishability of the blocks $w_{5,l}$ and $w_{4,l}$ from $w_{3,l}$, $l \geq 1$, we have $\eta_1^i = \beta_1^2$ for $i = 2, 5, \dots$ and $\varepsilon_1^i = \beta_1^2$ for $i = 2, 5, \dots$. Moreover, since the vectors $w_{4,l}$

and $w_{5,l}$, $l \geq 1$, are distinguishable, we have $\eta_1^i = \beta_1^3$ for $i = 3, 6, \dots$ or $\varepsilon_1^i = \beta_1^4$ for $i = 3, 6, \dots$. If now the second possibility happens, i.e. $\varepsilon_1^i = \beta_1^4$ for $j = 3, 6, \dots$, then we would change the order of the coordinates applying the permutation $(2\ 3\ 4)$. If the first possibility takes place, i.e. $\eta_1^i = \beta_1^3$ for $i = 3, 6, \dots$, then we use the permutation $(2\ 3\ 4\ 5)$. Consequently, we obtain

$$\begin{aligned} w_{2,l} : & \quad a_1 \quad \beta_l^1 \quad \gamma_1^{3l-5} \quad \beta_1^3 \quad \beta_1^4 \quad a_2, \\ w_{3,l-1} : & \quad a_1 \quad \delta_1^{3l-5} \quad \beta_l^2 \quad \delta_1^{3l-4} \quad \beta_1^4 \quad a_2, \\ w_{4,l-1} : & \quad a_1 \quad \varepsilon_1^{3l-5} \quad \varepsilon_1^{3l-4} \quad \beta_l^3 \quad \varepsilon_1^{3l-3} \quad a_2, \\ w_{5,l-1} : & \quad a_1 \quad \eta_1^{3l-5} \quad \eta_1^{3l-4} \quad \eta_1^{3l-3} \quad \beta_l^4 \quad a_2, \quad l \geq 2. \end{aligned}$$

If $\gamma_1^i = \beta_1^2$ for $i = 1, 4, \dots$, then the vectors $w_{2,1}$, $w_{2,l}$, $l \geq 2$, determine a column. Therefore, we can assume that at least one of the coordinates γ_1^i , $i = 1, 4, \dots$, is distinct from β_1^2 . We can suppose that it is γ_1^1 . Then, by the distinguishability of the vectors $w_{2,2}$ and $w_{3,l}$, $l \geq 1$, we have $\delta_1^i = \beta_1^1$ for $i = 1, 4, \dots$. If now $\delta_1^i = \beta_1^3$ for $i = 2, 5, \dots$, then the vectors $w_{2,1}$, $w_{3,l}$, $l \geq 1$, determine a column. Therefore, we can assume that $\delta_1^2 \neq \beta_1^3$. Now, let us take into account the vectors $w_{2,1}$, $w_{2,2}$ and $w_{3,1}$. Consequently, they have the following form

$$\begin{aligned} w_{2,1} : & \quad a_1 \quad \beta_1^1 \quad \beta_1^2 \quad \beta_1^3 \quad \beta_1^4 \quad a_2, \\ w_{2,2} : & \quad a_1 \quad \beta_2^1 \quad \gamma_1^1 \quad \beta_1^3 \quad \beta_1^4 \quad a_2, \\ w_{3,1} : & \quad a_1 \quad \beta_1^1 \quad \beta_2^2 \quad \delta_1^2 \quad \beta_1^4 \quad a_2. \end{aligned}$$

Passing to an appropriate isomorphic system, in much the same way as done before, we can assume that the latter system of vectors has the form

$$\begin{aligned} w_{2,1} : & \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1 \quad a_1, \\ w_{2,2} : & \quad a_1 \quad a_2 \quad \gamma_1^1 \quad a_1 \quad a_1 \quad a_1, \\ w_{3,1} : & \quad a_1 \quad a_1 \quad a_2 \quad \delta_1^2 \quad a_1 \quad a_1. \end{aligned}$$

Let us consider all vectors with codes $(1, 1, 2, l, 1, 1)$, $l \geq 2$. By their distinguishability from $w_{2,1}$, $w_{2,2}$ and $w_{3,1}$, they are as follows

$$w_{6,l-1} : \quad \rho_1^{3l-5} \quad a_1 \quad a_2 \quad \delta_l^2 \quad \rho_1^{3l-4} \quad \rho_1^{3l-3}, \quad l \geq 2.$$

If $\rho_1^i = a_1$ for all $i \geq 1$, then the vectors $w_{3,1}, w_{6,l}, l \geq 1$, determine a column. Suppose that at least one of the coordinates $\rho_1^i, i \geq 1$, is distinct from a_1 . We can assume that it is ρ_1^1 . (If not, then we would change the order of coordinates and the code ε_4 appropriately.) Take all vectors with codes $(l, 1, 2, 2, 1, 1), l \geq 2$. They can be written as follows

$$w_{7,l-1} : \begin{matrix} \rho_1^1 & a_1 & a_2 & \delta_2^2 & \sigma_1^{2l-3} & \sigma_1^{2l-2}, \end{matrix} \quad l \geq 2.$$

If $\sigma_1^i = \rho_1^2$ for $i = 1, 3, \dots$ and $\sigma_1^i = \rho_1^3$ for $i = 2, 4, \dots$, then the vectors $w_{6,1}, w_{7,l}, l \geq 1$, determine a column. Assume that $\sigma_1^1 \neq \rho_1^2$. (If not, then we would change the order of coordinates and the code ε_1 in an appropriate way.) Then we can also assume that $\rho_1^2 \neq a_1$. (If $\rho_1^2 = a_1$, and $\sigma_1^1 \neq a_1$, then we would replace the code ε_1 by ε'_1 so that $\varepsilon'_1 = (1\ 2) \circ \varepsilon_1$ on $\rho_1^1 + \mathbb{Z}$ and $\varepsilon'_1 = \varepsilon_1$ on the complement of $\rho_1^1 + \mathbb{Z}$.) Consider all vectors with codes $(1, 1, 2, 2, l, 1), l \geq 2$. They have the form

$$w_{8,l-1} : \begin{matrix} \tau_1^{2l-3} & a_1 & a_2 & \delta_2^2 & \rho_l^2 & \tau_1^{2l-2}, \end{matrix} \quad l \geq 2.$$

Since $\sigma_1^1 \neq \rho_1^2$, by the distinguishability of $w_{7,1}$ and $w_{8,l}, l \geq 1$, we have $\tau_1^i = \rho_1^1$ for $i = 1, 3, \dots$. If now $\tau_1^i = \rho_1^3$ for $i = 2, 4, \dots$, then the vectors $w_{6,1}, w_{8,l}, l \geq 1$, determine a column. Therefore, let us assume that $\tau_1^2 \neq \rho_1^3$ and $\rho_1^3 \neq a_1$. (If not, then we would change the code ε_5 in an appropriate way.) Take all vectors with codes $(1, 1, 2, 2, 1, l), l \geq 2$. They are as follows

$$w_{9,l-1} : \begin{matrix} \xi_1^{2l-3} & a_1 & a_2 & \delta_2^2 & \xi_1^{2l-2} & \rho_l^3, \end{matrix} \quad l \geq 2.$$

The distinguishability of $w_{8,1}$ and $w_{9,l}, l \geq 1$, implies $\xi_1^i = \rho_1^2$ for $i = 2, 4, \dots$, as $\tau_1^2 \neq \rho_1^3$. Since $w_{7,1}$ and $w_{9,l}, l \geq 1$, are distinguishable, we have $\xi_1^i = \rho_1^1$ for $i = 1, 3, \dots$ or $\sigma_1^i = \rho_1^3$ for $i = 2, 4, \dots$. If the first possibility happens, then the vectors $w_{6,1}, w_{9,l}, l \geq 1$, determine a column. Hence $\sigma_1^i = \rho_1^3$ for $i = 2, 4, \dots$. As a result we obtain

$$w_{6,1} : \begin{matrix} \rho_1^1 & a_1 & a_2 & \delta_2^2 & \rho_1^2 & \rho_1^3, \end{matrix}$$

$$w_{7,1} : \begin{matrix} \rho_2^1 & a_1 & a_2 & \delta_2^2 & \sigma_1^1 & \rho_1^3, \end{matrix}$$

$$w_{8,1} : \begin{matrix} \rho_1^1 & a_1 & a_2 & \delta_2^2 & \rho_2^2 & \tau_1^2, \end{matrix}$$

$$w_{9,1} : \begin{matrix} \xi_1^1 & a_1 & a_2 & \delta_2^2 & \rho_1^2 & \rho_2^3. \end{matrix}$$

Passing to an appropriate isomorphic system, in much the same way as done

before, we can assume that the latter system of vectors has the form

$$w_{6,1} : a_1 \ a_1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{7,1} : a_2 \ \sigma_1^1 \ a_1 \ a_1 \ a_1 \ a_1,$$

$$w_{8,1} : a_1 \ a_2 \ \tau_1^2 \ a_1 \ a_1 \ a_1,$$

$$w_{9,1} : \xi_1^1 \ a_1 \ a_2 \ a_1 \ a_1 \ a_1.$$

By Lemma 1, the set T contains vectors determining a column. □

References

- [1] G. HAJÓS, Über einfache und mehrfache Bedeckung des n -dimensionalen Raumes mit einem Würfelgitter, *Math. Z.* **47** (1941), 427–467.
- [2] O.H. KELLER, Über die lückenlose Erfüllung des Raumes Würfeln, *J. Reine Angew. Math.* **163** (1930), 231–248.
- [3] O.H. KELLER, Ein Satz über die lückenlose Erfüllung des 5- und 6-dimensionalen Raumes mit Würfeln, *J. Reine Angew. Math.* **177** (1937), 61–64.
- [4] J. C. LAGARIAS and P. W. SHOR, Keller's cube-tiling conjecture is false in high dimensions, *Bull. Amer. Math. Soc.* **27** (1992), 279–287.
- [5] J. MACKEY, A cube tiling of dimension eight with no facesharing, *Discr. Comput. Geom.* **28** (2002), 275–279.
- [6] H. MINKOWSKI, *Diophantische Approximationen*, Teubner, Leipzig, 1907.
- [7] O. PERRON, Über lückenlose Ausfüllung des n -dimensionalen Raumes durch kongruente Würfel I, II, *Math. Z.* **46** (1940), 1–26, 161–180.